

SweetPro®

CattleKandi

Supplement Block for Starting

Calves, Receiving Cattle, Stockers & Cows on Pasture

Net Wt. 102 kg.
(225 lbs.)

Reg. No. 621312

Guaranteed Analysis

This feed contains added selenium at 5 mg/kg

Crude Protein, Min.....	18.00%	Potassium (K), Actual.....	1.10%
Crude Fat, Min.....	5.00%	Magnesium (Mg), Actual.....	2.00%
Crude Fiber, Max.....	5.50%	Flourine (F), Max.....	0.56 mg/kg
Calcium (Ca), Actual.....	1.50%	Cobalt (Co), Actual.....	4.0 mg/kg
Phosphorus (P), Actual.....	1.00%	Copper (Cu), Actual.....	600 mg/kg
Salt (NaCl), Actual.....	7.75%	Iodine (I), Actual.....	8.0 mg/kg
Sodium (Na), Actual.....	3.05%	Iron (Fe), Actual.....	400 mg/kg
Sulfur (S), Actual.....	0.40%	Manganese (Mn), Actual.....	750 mg/kg
		Zinc (Zn), Actual.....	1,200 mg/kg
Vitamin A, Minimum.....	220,000 IU/kg		
Vitamin D, Minimum.....	44,000 IU/kg		
Vitamin E, Minimum.....	180 IU/kg		

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Corn Gluten Meal, Magnesium Oxide, Monocalcium Phosphate, Potassium Chloride, Calcium Carbonate, Canola Oil, Yeast Culture Dehydrated, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Protein-ate (Reg. 990743), Manganese Protein-ate (Reg. 990744), Aspergillus Oryzae Extract Dehydrated, Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Acetate, Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018).



Cattle Kandi

Directions for Use

1. Before feeding SweetPro® Cattle Kandi lick blocks for the first time, animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Have one lick block available per 25 head of cattle.
4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
5. Place blocks near water sources or areas cattle frequent regularly.
6. Make blocks available in more than one location in large pastures or where more than one water source is available.
7. If animals run out of lick blocks, repeat from Step#1.
8. Other salt and minerals may be provided free choice to balance specific needs.
9. A two week adaptation period will be required to establish a base consumption rate.
After adaptation period, recommended range of intake is 0.34 to 0.56 kgs. per day. Calves .1 to .5 kg/day.
10. Once animals consistently consume more than 0.56 kgs/hd/day, transition to SweetPro 16. If daily consumption is consistently below 0.34 kgs/hd/day, transition to SweetPro® Starter.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**
Do not use in association with another feed containing supplemental selenium.

Control #:

Registered by Harvest Fuel, Inc. dba SweetPro Feeds, Walhalla, ND 58282 (800) 327-9222

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. Made in U.S.A. Printed with soy inks.
Patent NO.: US 5,260,089 Patent NO.: US 6,777,019 B1 Patent NO.: US D352,136 Patent NO.: US D440,362 Patent NO.: US D428,272