MUZZEJI (114 kg.) **Net Wt. 250 lbs.**

ENCY

Guaranteed Analysis

	Minimum 20.00%
(includes not more that	an 8% equivalent protein from non-protein-nitrogen)
Crude Fat	
Calcium (Ca)	
Salt (NaCl)	
Magnesium (Mg)	
Potassium (K)	
Sulfur (S)	
Copper (Cu)	Minimum 600 ppn
Selenium (Se)	Minimum 4.75 ppm - Maximum 5.25 ppn
Zinc (Zn)	Minimum 1,200 ppn
Manganese (Mn)	
Cobalt (Co)	Minimum 4.0 ppn
Iodine (I)	
Vitamin A	
Vitamin D	
Vitamin E	
Alpha-Amylase	Minimum 3,393 ug starch hydrolyzed/min/oz
Amyloglucosidase	Minimum 513 ug maltose hydrolyzed/min/oz

Ingredients: Corn Condensed Distillers Solubles, Corn Distillers Dried Grains with Solubles, Salt, Fermented Corn Protein, Magnesium Oxide, Mono-Calcium Phosphate, Yeast Culture, Wheat Starch, Vegetable Oil (Canola and Flax Oil), Calcium Carbonate, Ammonium Sulfate, Biuret, Calcium Sulfate, Citric Acid, Potassium Chloride, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Dried Aspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

FiberMate™ 20

SweetPro® FiberMate 20 lick blocks are designed for essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. FiberMate 20 should be used for cattle on average to below average forages. Daily intake will vary depending on forage quality and 🔲🖼 nutritional needs.



Feeding Instructions:

- 1. Before feeding SweetPro® FiberMate 20 lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Provide an adequate source of good quality roughage and forage at all times (all
- 4. Have one lick block available per 25 head of beef or dairy cattle.
- 5. Place lick blocks near water sources or areas cattle frequent regularly.
- 6. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- 8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs./hd/
- 9. Once animals consistently consume more than 1.25 lbs./hd/day, transition to SweetPro®Super25. If daily consumption is consistently below .75 lbs./hd/day, transition to SweetPro® FiberMate 18.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 600 ppm copper. DO NOT FEED TO SHEEP!

Do not use in association with another feed containing supplemental selenium.





