Suczt Dro[®] Net Wt. 250 lbs.

.Minimum 3.393 ug starch hydrolyzed/min/oz.

..Minimum 513 ug maltose hydrolyzed/min/oz.

Guaranteed Analysis	
Crude Protein	
(includes not more than 15% equivalent protein from non-protein-nitrogen)	
Crude Fiber	
Calcium (Ca)	
Phosphorus (P)	
Salt (NaCl)	
Magnesium (Mg)	
Potassium (K)	
Sulfur (S)	
	Minimum 600 ppm
Manganese (Mn)	
	Minimum 4.0 ppm
	Minimum 80 IU/lb.

Ingredients: Corn Condensed Distillers Solubles, Corn Distillers Dried Grains with Solubles, Salt, Ammonium Sulfate, Fermented Corn Protein, Magnesium Oxide, Calcium Carbonate, Mono-Calcium Phosphate, Corn Gluten Meal, Biuret, Yeast Culture, Calcium Sulfate, Vegetable Oil (Canola and Flax Oil), Potassium Chloride, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Magnum™

SweetPro® Magnum™ blocks are designed for essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Magnum should be used for cattle on below average to poor and harsh forages. Daily intake will vary depending on forage quality and nutritional needs.



Feeding Instructions:

- Before feeding SweetPro® Magnum lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 4. Have one lick block available per 25 head of beef cattle.
- 5. Place blocks near water sources or areas cattle frequent regularly.
- Make blocks available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs. per day.
- 9. Once animals consistently consume less than .75 lbs./hd/day, transition to SweetPro $\!\!^{\otimes}$ Super 25.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #:



Alpha-Amylase

Amyloglucosidase

